

## Exotic hadron holography from anomalous dimensions

---

**Hilmar Forkel\***

*Institut für Physik, Humboldt-Universität zu Berlin, D-12489 Berlin, Germany*

*E-mail: forkel@physik.hu-berlin.de*

The anomalous dimensions of hadronic interpolators contain dynamical information on the properties of the associated hadron states. We point out that they provide, in particular, a link by which gauge-invariant information on exotic contributions to hadronic wavefunctionals can be obtained from approximate gravity duals for QCD. This is demonstrated by the holographic description of a dominant tetraquark component in the lightest scalar mesons.

*Xth Quark Confinement and the Hadron Spectrum,*

*October 8-12, 2012*

*TUM Campus Garching, Munich, Germany*

---

\*Speaker.

While the anomalous dimensions of color-singlet operators play a central role in the original, conformal versions of the gauge/string correspondence [1], they have only recently begun to enter holographic approaches to QCD. In particular, anomalous dimensions of hadronic interpolators were implemented to provide an AdS/QCD [2] description of quark correlations inside hadrons [3, 4] which can have a significant and in exotic cases even striking impact on the hadron properties. Although multi-quark components in hadronic wave functionals are typically gauge dependent, a holographic description is still possible because the five-dimensional bulk modes are dual to the interpolators of the corresponding hadrons. Hence the anomalous dimensions of these interpolators import gauge-invariant information on their quark content and couplings, and thus on the multi-quark correlations in the corresponding hadron, as bulk-mode mass corrections into the gravity dual.

This AdS/QCD representation of multi-quark effects was originally introduced to describe di-quark correlations in baryons [3]. It changes the resulting light-quark baryon excitation spectrum into

$$M_{n,L}^2 = 4\lambda^2 \left( n + L + \frac{3}{2} \right) - 2(M_\Delta^2 - M_N^2) \kappa \quad (1)$$

(where  $\lambda$  is the IR scale of the “metric soft-wall” gravity dual [5] while  $n$  ( $L$ ) denotes the radial (angular momentum) excitation level). The second term, proportional to the baryon’s “good-diquark fraction”  $\kappa$ , is generated by suitable anomalous dimensions for the QCD nucleon interpolators. Equation (1) describes the linear square-mass trajectories of the over 40 measured nucleon and delta (with  $\kappa = 0$ ) resonances with unprecedented accuracy. The dual mode solutions further reveal that baryons with larger  $\kappa$  have a smaller size.

Encouraged by these results, the anomalous-dimension-induced representation of multi-quark correlations was then applied to the more challenging holographic description of exotic hadrons with a non-standard (valence) quark content. The light scalar meson sector [6] with its expected tetraquark component [7] was examined in Ref. [4]. The radial bulk equation for the modes dual to the scalars can be written as the Sturm-Liouville problem  $[-\partial_z^2 + V(z)] \phi(q, z) = q^2 \phi(q, z)$ . In the dilaton soft-wall gravity dual [8] without anomalous-dimension contributions, the potential  $V$  has the form

$$V(z) = \left( \frac{15}{4} + m_5^2 R^2 \right) \frac{1}{z^2} + \lambda^2 (\lambda^2 z^2 + 2). \quad (2)$$

The anomalous dimension  $\gamma(z)$  of the tetraquark interpolator  $J_{\bar{q}^2 q^2}$  (i.e. the local four-quark operator which most strongly couples to the tetraquark state) with scaling dimension  $\Delta_{\bar{q}^2 q^2} = 6 + \gamma(z)$  adds the universal contribution

$$\Delta V(z) = \gamma(z) [\gamma(z) + 8] \frac{1}{z^2} \quad (3)$$

to the potential (2) with  $m_5^2 R^2 = 12$ . Eq. (3) implies the crucial lower bound  $\Delta V(z) \geq -16/z^2$  which holds for any  $\gamma$  and prevents the collapse of the dual modes into the AdS<sub>5</sub> boundary. This bound is saturated by  $\gamma \equiv -4$  and therefore determines the lightest tetraquark mass

$$M_{\bar{q}^2 q^2, 0} \geq M_{\Delta=2, 0} = 2\lambda \quad (4)$$

which the anomalous-dimension-induced holographic binding mechanism can produce. Moreover, for constant values  $-4 < \gamma < -3$  the tetraquark ground state is lighter than its  $\bar{q}q$  counterpart. Since

$\gamma$  only enters through the mass term of the bulk mode which is model-independently prescribed by the AdS/CFT dictionary, the correction (3) and the associated binding mechanism will arise in other AdS/QCD duals as well.

To estimate the quantitative impact of the anomalous-dimension contribution  $\Delta V$  (until direct QCD information on the RG flow of  $\gamma$  will eventually become available and fix  $\Delta V$  uniquely), a typical power ansatz  $\gamma(z) = -az^\eta + bz^\kappa$  can be adopted. Its coefficients turn out to be tightly constrained by consistency and stability requirements but can still produce almost maximal ground-state binding [4]. The latter drives the mass  $M_{\bar{q}^2 q^2, 0}$  of the lightest tetraquark from  $\sim 40\%$  above (for  $\gamma \equiv 0$ ) down to  $\sim 20\%$  below the  $\bar{q}q$  ground-state mass  $M_{\bar{q}q, 0} = \sqrt{6}\lambda$ . The resulting masses  $M_{\bar{q}^2 q^2, n}$  of the tetraquark excitations get pushed beyond the corresponding  $M_{\bar{q}q, n}$  from around  $n \gtrsim 2$ . The higher-lying radial tetraquark excitations will therefore likely be broad enough to prevent the appearance of supernumerary states in the scalar meson spectrum.

It should be interesting to extend the anomalous-dimension-based holographic description of non-valence quark components to other exotics, including heavy tetraquarks, pentaquarks and hybrids. Moreover, anomalous-dimension-induced corrections also encode other aspects of hadronic structure which largely remain to be explored.

## Acknowledgments

It is a pleasure to thank the organizers for a very informative and enjoyable conference.

## References

- [1] O. Aharony et al., *Large- $N$  field theories, string theory and gravity*, Phys. Rep. **323** (2000) 183.
- [2] Y. Kim and D. Yi, *Holography at work for nuclear and hadron physics*, Adv. High Energy Phys. **2011** (2011) 259025; S.J. Brodsky and G.F. de Téramond, *AdS/CFT and Light-Front QCD*, arXiv:0802.0514.
- [3] H. Forkel and E. Klempt, *Di-quark correlations in baryon spectroscopy and holographic QCD*, Phys. Lett. **B 679** (2009) 77.
- [4] H. Forkel, *Light scalar tetraquarks from a holographic perspective*, Phys. Lett. **B 694** (2010) 252; *Multiquark correlations in light mesons and baryons from holographic QCD*, AIP Conf. Proc. **1388** (2011) 182 [arXiv:1103.3902].
- [5] H. Forkel, M. Beyer and T. Frederico, *Linear square-mass trajectories of radially and orbitally excited hadrons in holographic QCD*, JHEP **07** (2007) 077; *Linear meson and baryon trajectories in AdS/QCD*, Intl. J.Mod. Phys. E **16** (2007) 2794.
- [6] E. Klempt and A. Zaitsev, *Glueballs, hybrids, multiquarks – Experimental facts versus QCD inspired concepts*, Phys. Rep. **454**, 1 (2007); C. Amsler and N.A. Törnqvist, *Mesons beyond the naive quark model*, Phys. Rep. **389**, 61 (2004); D.V. Bugg, *Four sorts of meson*, Phys. Rept. **397**, 257 (2004); E. Ruiz Arriola and W. Broniowski, *Scalar-isoscalar states in the large- $N_c$  Regge approach*, Phys. Rev. D. **81** (2010) 054009.
- [7] R.L. Jaffe, *Phenomenology of  $Q^2 \bar{Q}^2$  mesons*, Phys. Rev. D **15**, 267, 281 (1977); J.R. Peláez, *Nature of light scalar mesons from their large- $N_c$  behavior*, Phys. Rev. Lett. **92** (2004) 102001.
- [8] A. Karch, E. Katz, D.T. Son and M.A. Stephanov, *Linear Confinement and AdS/QCD*, Phys. Rev. D **74** (2006) 015005.